



# Hydrothermal alteration in the NW Hellas Region

Eldar Noe Dobrea

Gregg Swayze

NOTE ADDED BY JPL WEBMASTER: This content has not been approved or adopted by, NASA, JPL, or the California Institute of Technology. This document is being made available for information purposes only, and any views and opinions expressed herein do not necessarily state or reflect those of NASA, JPL, or the California Institute of Technology

0 320 km

# Hellas

- Formed in Early Noachian by impact of ~250 km object (*e.g.*, Leonard and Tanaka, 2001)
- Subsequent modification by aeolian, fluvial, glacial/periglacial, volcanic processes gave rise to a diversity of geological environments and landforms inside Hellas and in the circum-Hellas region.
- Of particular interest to the question of *habitability* is the possibility that a variety of aqueous systems may have developed in the region after the Hellas impact.
  - Hydrothermal systems (*e.g.*, Newsom, 1980).
  - Marine/lacustrine (*e.g.*, Wilson *et al.*, 2007; Condit *et al.*, 2010 and references therein)
- Energy deposited by impact should have produced elevated temperatures for a prolonged period (how long? how hot?)
- Impact also thought to have triggered vent volcanism (*e.g.*, Williams *et al.*, 2009)
- Evidence for aqueous activity in the region during that time period:
  - Mineralogical: phyllosilicates in Tyrrhena Terra (Pelkey *et al.*, 2007; Loizeau *et al.*, 2009) and around Hellas (*e.g.*, Ansan *et al.*, 2011; Crown *et al.*, LPSC 2011).
  - Morphological: heavy dissection of the region, layered intracrater deposits




Primary mineral or rock	Reported alteration minerals	Hydrothermal Alteration environment	Reference
Basalt	Kaolinite, smectite, jarosite, alunite	Acidic pH	Morris <i>et al.</i> , 2001; Morris <i>et al.</i> , 2003
Pyroxene-amphibole andesite	Cristobalite, alunite, pyrite, kaolinite, goethite, hematite	Acidic pH	Isobe and Korenaga, 2010
Basalt	Mg-carbonate, talc	Neutral to basic pH	Brown <i>et al.</i> , 2010
Wollastonite	Mg-montmorillonite, talc, mixed layer stevensite/chlorite	Neutral to basic pH groundwater	DeRudder and Beck, 1963; Whitney and Eberl, 1982
Granite, K-feldspar	Kaolinite, muscovite, biotite, halloysite	-	Thomas and Walter, 2004;
Impact melt rock	Fe-chlorite, Fe smectite, silica, K feldspar, zeolite	-	Newsom, 1980; Allen <i>et al.</i> , 1982
Olivine	serpentine	-	Normand <i>et al.</i> , 2002
Basalt, grabbro	Prehnite, quartz, calcite, epidote	Also in low-grade metamorphism	Freedman <i>et al.</i> , 2009

# Potential Hydrothermal/low grade metamorphic alteration products on Mars

- Prehnite: Nili Fossae (Ehlman *et al.*, 2009), Argyre rim (Buczowski *et al.*, 2010), NW Hellas (Crown *et al.*, LPSC 2011)
- Serpentine: Nili Fossae (Ehlman *et al.*, 2009)
- Chlorite: Nili Fossae (*e.g.*, Ehlman *et al.*, 2009; Poulet *et al.*, 2005); NW Hellas (Crown *et al.*, LPSC 2011)
- Hydrated silica: Nili Fossae (Skok *et al.*, this session)
- Of particular interest is Prehnite ( $\text{Ca}_2\text{Al}_2\text{Si}_3\text{O}_{10}(\text{OH})_2$ )
  - Metamorphic grade transitional between zeolite and greenschist facies
  - Forms under specific conditions: 2-7 kbar, 200 – 350°C, and  $X_{\text{CO}_2} < 0.004$  (Blatt and Tracy, 1995)
  - Typically associated with chlorite and pumpellyite (*e.g.*, Frey and Robinson, 1999).


# Results




- Prehnite:  
absorptions at 2.35-2.36, 1.48, 2.23, 2.28, 2.57  $\mu\text{m}$
- Chlorite:  
absorptions at 1.40, shoulder at 2.25-2.26, 2.33-2.35
- Serpentine: absorptions at 1.40, 2.32-2.33, 2.50-2.51
- “Vermiculite” – mixed layer vermiculite/biotite.  
Can also be other mixed layer smectite/chlorite: absorption at 1.92, steep drop-off at 2.30-2.31
- Illite/muscovite -  
absorptions at 2.2, 2.35

# Results

Spectral library hydrothermal alteration products



Examples of alteration products in NW Hellas observed in CRISM data






# Geological Setting



NW Hellas Region preserves record of diverse geological processes

0 320 km


A horizontal scale bar with vertical tick marks. The number '0' is at the left end and '320 km' is at the right end.



Dissected units above 1000 m

Heavy dissection

MOLA Elevation (m)



0 500 60 km 500



# NW Hellas Region

N




Preserves record of diverse geological processes

0


320 km


















Prehnite units

N




BD1900R



High : 255

Low : 0

0 1.5 3 6 km






# FRT00019C80


N




Prehnite (ripple-forming unit)



Prehnite + Chlorite  
(units contains ripple-  
forming material and  
indurated rough unit)



**BD1900R**




0 0.35 0.7 1.4 km



FRT0001214D

N



Illite/muscovite  
(rocky unit)

Prehnite

Prehnite + Chlorite  
(Rough rocky unit)




0 0.45 0.9 1.8 km



FRT0001214D

Prehnite  
(smooth and  
ripple-forming  
unit)


BD1900R



0 0.15 0.3 0.6 km









# Pitted Plains Deposits



Yardanged  
Layered  
100-250 meters deep  
Occur at elevations  $< -500$  m

0 0.8 km

A horizontal scale bar with vertical tick marks. The text '0' is at the left end and '0.8 km' is at the right end.



# Pitted Crater Floors


Polygonally-fractured unit

Scooped unit

Layers

Layered  
Morphologic stratigraphy  
~300-500 m thick

0 0.18 km







# Pitted Plains Deposits: FRT0001366A

N



Mixed-layer S/C →



D2300



High : 255

Low : 0

0

4 km



# Geological inferences

- We observe a variety of alteration products including prehnite, chlorite, illite/muscovite, mixed-layer S/C or B/V, saponite.
- Prehnite+chlorite+illite in close spatial association on plains unit strongly suggestive of hydrothermal activity.
  - observed in Hilly Unit, which has been interpreted to be uplifted crustal material and ejecta from Hellas impact and post-impact structural and erosional modifications.
- Subsequent erosion of Noachian plains material resulted in dissection and redeposition to form smooth embaying plains and intracrater units 200-400 m in thickness.
- Low-grade metamorphic materials (mixed-layer S/C) identified within pit walls in plains units.
- Plains units have very gentle ( $<2.5^\circ$ ) slopes throughout, and are easily accessible to landing system. Scarcity of boulders in this unit would make traversability very straightforward.







- Unit above 500 m – Npl2 – Subdued cratered unit – Forms widespread moderate to heavily cratered, relatively smooth plains marked by subdued crater rims, small channels, ridges, and uneven terrain. Crater floors partly to completely infilled with smooth material; ejecta blankets rare. Some heavily eroded craters dissected by small channels. Faults rare. Material gradational with most adjacent units. *Interpretation* – Ancient veneer of aeolian, fluvial, and perhaps volcanic materials that partly resurface underlying cratered and dissected units (units Npl1 and Npld).
- Prehnite in unit Nh – Hilly Unit – Forms rugged, high-relief, densely cratered terrain with numerous isolated massifs. Channels and ridges common; faults along west rim. *Interpretation*: Uplifted crustal material and ejecta from Hellas impact and post-impact structural and erosional modifications.
- Smooth pitted plains in unit Hpl3 – Smooth unit – Forms moderately cratered, smooth, flat to undulating, relatively featureless plains and patches around the rim and within highlands surrounding Hellas basin. Channel common; faults and flow fronts rare. Embays all other materials of plateau sequence and fills many impact craters. *Interpretation*: Thick fluvial, aeolian, and volcanic deposits burying most underlying rocks
- (Leonard and Tanaka, 2001)



# Massifs: FRT00013D40




Saponite

Mixed-layer

Knob exposed in pitted  
plains unit  
Mixed layer at base  
Saponite at top  
Top of knob rises above  
surrounding pitted plains

D2300



0 2 km